

First-principles Design of Nanomachines

Jayanth R. Banavar,¹ Marek Cieplak,² Trinh Xuan Hoang,^{1,3} and Amos Maritan⁴

¹*Physics Department, Penn State University, 104 Davey Lab, University Park, PA 16802, USA*

²*Institute of Physics, Polish Academy of Sciences, Aleja Lotnikow 32/46, 02-668 Warsaw, Poland*

³*Institute of Physics, Vietnamese Academy of Science and Technology, 10 Dao Tan, Ba Dinh, Hanoi 10000, Vietnam*

⁴*Dipartimento di Fisica ‘G. Galilei’, Università di Padova & CNISM,
unità di Padova & INFN, sezione di Padova, Via Marzolo 8, 35131 Padova, Italy*

Learning from nature’s amazing molecular machines, globular proteins, we present a framework for the predictive design of nano-machines. We show that the crucial ingredients for a chain molecule to behave as a machine are its inherent anisotropy and the coupling between the local Frenet coordinate reference frames of nearby monomers. We demonstrate that, even in the absence of heterogeneity, protein-like behavior is obtained for a simple chain molecule made up of just thirty hard spheres. This chain spontaneously switches between two distinct geometries, a single helix and a dual helix, merely due to thermal fluctuations.

Significant advances in laboratory techniques for tailoring and processing materials at the atomic scale have resulted in nanotechnology becoming an increasingly mature field with great promise. One of the exciting goals of the field is the design of powerful machines, such as functional entities that can switch reversibly between distinct geometries^{1,2,3,4}. Such machines would not only be of great use on their own but also could yield novel emergent behavior on networking them together⁵. The existence of life in its myriad forms provides a proof-of-concept of what one might aspire to accomplish with nanotechnology.

Proteins are complex water-soluble chain molecules made up of tens or hundreds of twenty types of naturally occurring amino acids and exhibit conformational switching⁶ triggered by influences such as ligand binding. At the nanoscale, thermal fluctuations yield forces with magnitude comparable to those involved in chemical reactions catalyzed by the proteins⁷. How might one design a machine whose functionality is structure-based? In its simplest form, we seek an object which takes on just a few distinct geometries in a reproducible manner and is able to switch reversibly between them due to thermal fluctuations. Such a situation would allow external stabilizing influences to favor a given conformation over the others and allow for the development of powerful machines at the nanoscale.

A collection of hard spheres constitutes the simplest model of matter and exhibits both a crystalline phase and a fluid phase on varying the density of spheres⁸. A linear chain of hard spheres is the simplest connected object with the fewest constraints and thus the greatest flexibility⁹. Such a chain, at high temperatures or when there are no interactions promoting compaction, would occupy a random coil phase in which all self-avoiding conformations are equally likely. This situation is not conducive for machine design. In the presence of inter-sphere interactions promoting compaction and when there are no frustrating influences from the sphere tethering, one would expect a generic compact phase in which, at least locally, the preferred conformation is that of a face-centered cubic (fcc) lattice. It

was conjectured by Kepler, and proven more recently by Hales, that the fcc structure provides for optimal packing of unconstrained spheres¹⁰. There is again a high degeneracy of compact conformations with no assurance that any given conformation will be reached from a random coil conformation rapidly and reproducibly. Again, this situation is not conducive for the design of machines.

Of course, what one requires is a phase of matter with much fewer ground state conformations than either the random coil or generic compact phase. Additionally, one needs a phase which is in the vicinity of a phase transition (or crossover for finite size systems) to a distinct phase, because that provides for exquisite sensitivity to the right types of external influences. Our thinking is guided by the liquid crystal phase⁸ which is a distinct state of matter that is poised in the vicinity of the liquid phase. This phase is known to be one of the most sensitive phases of matter. The liquid crystal phase opens up because of the anisotropy of the constituent molecules – there is no longer a need for simultaneous ordering in all three directions. Rather, one can have translational order in fewer than three dimensions and orientational order. Here we address the issue of how one might open up a distinct phase with relatively few ground state conformations in the vicinity of the random coil phase. Such a phase would then be the analog of a liquid crystal phase but this time for chain molecules.

Armed with insights from the liquid crystal phase and the behavior of proteins, we identify two mechanisms for thinning the number of candidate ground state structures and demonstrate that these are sufficient for the creation of the sought after behavior. One can define a Frenet reference frame (a local Cartesian coordinate system) at each location of the chain molecule comprised of the tangent, the normal, and the binormal as the orthogonal coordinate axes. These coordinate systems play a crucial role in at least two ways, as can be readily seen in the protein context: first, the chemistry of hydrogen bonds and other chemical features yield constraints on the relative orientations between the coordinate systems associated with pairs of amino acids in contact; second, the side-chains of amino acids are located in a specified direction




FIG. 1: Sketch of the chain molecule. The backbone is modeled as a chain of (blue) spheres of radius a with a separation along the chain $l = 3.8\text{\AA}$. The nearest neighbor spheres along the backbone are allowed to overlap with each other thereby overtly introducing uniaxial anisotropy. $\mathbf{t}_i$ and $\mathbf{n}_i$ are the tangent and normal vectors assigned to each sphere, i , except to those at the ends of the chain. Side-spheres (shown in pink) of radius b are attached to the backbone spheres in the negative normal direction. The side-spheres are not allowed to overlap with either the backbone spheres or with each other.

with respect to these coordinate systems - for example, the amino acid side chains are typically pointed approximately in the outward normal direction. Indeed, both the α -helix and the β -sheet allow for the placement of side-chains in a manner that avoids steric clashes.

The first mechanism follows from the observation that a model of a chain comprised of spheres is unable to capture the inherent anisotropy induced by the presence of the chain constraint - at each location of the chain, there is a tangent direction defined by the two adjacent objects. Thus the simplest model capturing the correct symmetry is one in which the constituent monomers are no longer **isotropic**. We therefore allow for the overlap of van der Waals spheres of adjacent monomers along the chain. Such an overlap overtly breaks the isotropy of the spheres and confers uniaxial anisotropy to the chain. In both the emergent building blocks of protein structures, α -helices and β -sheets, nearby segments are placed right up against each other and aligned parallel to each other, reflecting the anisotropy. In a protein, the distance between neighboring C^α atoms is $3.8\text{\AA}$ requiring that the radius of the backbone of an amino acid be larger than $1.9\text{\AA}$. This constraint is easily met by considering the van der Waals radii of the constituent atoms. For example, one would estimate¹¹ that the smallest amino acid, glycine,




FIG. 2: Phase diagram of chain molecules made up of spheres. The panels (a, b) show the phase diagrams at $T = 0$ in the $a - R_c$ plane for chains of length $N=12$ without side-spheres ($b = 0$) and with side-spheres ($b = 2.5\text{\AA}$) respectively. The intermediate compact phase arises on the edge of compaction of the chain molecule when R_c becomes sufficiently large to allow the attractive interaction to be effective. The vicinity of this phase to other phases (it is sandwiched between two distinct phases) confers sensitivity to structures in the intermediate compact phase. The other panels show typical ground state conformations: a random coil conformation obtained with $R_c < 2a$ (c), a globule conformation obtained with $a = 2.5\text{\AA}$, $R_c = 9\text{\AA}$ (d), a saddle conformation obtained with $a = 3.2\text{\AA}$, $R_c = 7\text{\AA}$ (e), several helix conformations with different pitch to radius ratios obtained for $a = 2.6\text{\AA}$, $R_c = 6.4\text{\AA}$ (f), $a = 3.2\text{\AA}$, $R_c = 8\text{\AA}$ (g), and $a = 3.4\text{\AA}$, $R_c = 9\text{\AA}$ (h). The helical and saddle ground state conformations in the model with no side-spheres are retained even when side-spheres are present ($b = 2.5\text{\AA}$). However, the side-spheres eliminate several random globule and random coil conformations thereby stabilizing the intermediate compact phase.

has a radius larger than $2.4\text{\AA}$. Second, we study the effect of attaching side-chains to the backbone monomer, e.g., in the negative normal direction. The self-avoidance of these side spheres with each other and with the backbone spheres results in an induced coupling between pairs of Frenet frames.

We find that each of these features results in the creation of a distinct intermediate phase for short chains sandwiched between the generic compact phase and the random coil phase. Furthermore, when the two features are combined, the phase is stabilized and occupies a larger region in parameter space. For our simplified model of a chain of backbone spheres (Fig. 1) with a sphere separation along the sequence of $3.8\text{\AA}$, we have three length parameters: the radius, a , of the backbone spheres, the radius, b , of the side-spheres, and the cut-off




FIG. 3: Multiple specific heat peaks corresponding to three distinct phases. We study a chain of $N = 16$ with $a = 3\text{\AA}$, $b = 2.5\text{\AA}$, and $R_c = 7.5\text{\AA}$. (a) The temperature dependence of the specific heat indicates two peaks – one at $T \approx 0.2\epsilon/k_B$ and another at $T \approx 0.97\epsilon/k_B$. The lower temperature peak indicates a crossover into a single helix conformation from a dual helix, whereas the peak at the higher temperature indicates a crossover between the dual helix and the random coil phases. (b) Contour plot of the effective free energy as a function of energy and contact order, CO , at the lower transition temperature. CO is defined as the sum over sequence separation of all contacts divided by the product of the number of contacts and N . The effective free energy at a given temperature T is determined as $\mathcal{F}(E, CO) = -k_B T \ln P(E, CO)$, where $P(E, CO)$ is a weighted two-dimensional histogram for that temperature obtained using the multiple histogram method²⁵. The unweighted histograms at multiple temperatures are collected through parallel tempering²⁶ Monte Carlo simulations. The free energy difference between consecutive contour levels is 4ϵ . The contour plot shows two minima corresponding to the single helix and the dual helix as indicated.

scale, R_c , for the pairwise attractive interaction potential, ϵ/k_B , between the backbone spheres (See Materials and Methods). In the absence of side-spheres and for non-overlapping backbone spheres, $a \leq 1.9\text{\AA}$, there is a crossover from a random coil phase (in which the conformations are mostly extended) to a globule phase in which one has compact conformations with no distinct motifs such as helices. The situation becomes qualitatively different on incorporating one or both of the two key features.

Fig. 2a shows the $T = 0$ phase diagrams in the a - R_c

plane for a $N = 12$ chain. Distinct phases, corresponding to helical and saddle-like conformations, emerge between the coil and the globular phases analogous to the opening up of a liquid crystal phase between the liquid and crystalline phases for anisotropic molecules⁸. The saddle can be thought of as a piece of a dual helix structure and does develop into a dual helix for longer chains. (The dual helix is distinct from the double helix because the former is a conformation of a single chain whereas the latter is comprised of two chains.) These phases have a lower entropy than both the globule and random coil (swollen) phases but are stabilized by the attractive interaction potential. We also find that the presence of side-spheres even without overlap of the backbone spheres induces helical conformations (Figure S1 in SI). This effect is accentuated when both features are present simultaneously (Figure S2 in SI). Interestingly, conformations composed of nearly parallel strands similar to those in β -sheets can be obtained by adding a bending energy into the model (Figure S3 in SI).

We now consider an even shorter system ($N = 16$) with an enhanced overlap of the backbone spheres: $a = 3.0\text{\AA}$. The radius of the side-sphere is chosen to be $b = 2.5\text{\AA}$, and the range of attraction is now increased to $R_c = 7.5\text{\AA}$. These parameter values yield a helix ground state but is close to a crossover to the dual helix state. Fig. 3a shows that the specific heat has two peaks corresponding to two crossover temperatures, $T_1 \approx 0.2\epsilon/k_B$ and $T_2 \approx 0.97\epsilon/k_B$. At the lowest temperatures, the helix is the dominant conformation. At intermediate temperatures, one obtains the dual helix structure, whereas, at higher temperatures, one obtains a random coil. Fig. 3b shows the contour plots of the free energy on the energy-contact order plane at T_1 and exhibits pronounced minima corresponding to the single helix and dual helix. The helix is a bit lower in energy while the dual helix is entropically more favorable. The free energy barrier for the switching at T_1 is about 4ϵ which is equivalent to the energy required to break 4 contacts. This barrier is significantly higher than $k_B T_1$ and the two helical conformations are each quite stable at this temperature. Quite remarkably, our Monte Carlo simulations with standard pivot and crankshaft move sets show a dynamical switching between the two conformations in a single trajectory at a somewhat higher temperature of $T = 0.5\epsilon/k_B$. The system is bistable with a rapid switching between distinct conformations (for examples of molecular switches, see Refs.^{1,2,3,4}) with little weight for intermediate conformations (Fig. 4). Note that the switching can also be easily effected by means of an external influence which is sensitive to the chain end-to-end distance.

It is important to note that the ease with which one obtains this system without any fine-tuning of details is made possible by the existence of the intermediate phase. The conformations in this phase are in the vicinity of the random coil phase while retaining order due to the proximity of the compact phase. This results in special sensitivity to small perturbations induced, e.g., by ther-

mal fluctuations which, in the example presented here, are responsible for the switching between two distinct geometrical shapes. Unlike the random coil conformations which can switch from one to another easily due to thermal fluctuations, the structures in the intermediate phase exhibit some stability. At the same time, the structures are not so densely compact that they are subject to sluggish dynamics and kinetic inaccessibility characteristic of the glassy phase. These distinct advantages of the intermediate phase have a wider applicability than for chain molecules as evidenced by the sensitive liquid crystal phase sandwiched between the crystalline and liquid phases. In liquid crystals, the anisotropy arises from the asymmetric shape of the constituent molecules, whereas here the anisotropy is a natural consequence of the chain topology of the molecule. Interestingly, earlier computational studies^{13,14} had found that overlapping adjacent monomers yield helical conformations. Also protein-like folds are adopted by a host of non-biological polymers^{15,16,17,18,19,20,21} (see Appendix).

Understanding the properties of matter is greatly simplified on using the concept of its phases⁸. For example, a liquid possesses certain gross properties, such as adopting the shape of the container and its ability to flow, irrespective of its constituent molecules and their underlying chemistry. Globular proteins share a great deal of common characteristics - they are all linear chains of the same 20 amino acids, they fold rapidly and reproducibly into their native state structures, these structures are made up of emergent building blocks in the form of helices and almost planar sheets, the total number of distinct folds that proteins adopt is limited in number^{22,23} much as the number of space groups associated with Bravais lattices⁸ is 230, proteins are flexible and versatile in their folded state, and proteins have a tendency to aggregate and form amyloid which in turn is implicated in debilitating human diseases. These common attributes of proteins suggest that protein structures occupy an intermediate phase of matter^{12,24} which confers on them their many amazing characteristics.

Whitesides wrote⁵ that one ought to “take existing nanomachines—those present in the cell—and learn from them. We will undoubtedly be able to extract from these systems concepts and principles that will enable us to make variants of them that will serve our purposes, and others that will have entirely new functions”. The phase of matter so successfully used by nature as the basis of life is ready to be exploited in the laboratory.

Methods

We consider chains of N hard spheres each of radius a . The beads spacing along the chain is fixed to be $l = 3.8\text{\AA}$. Non-consecutive spheres are not allowed to overlap, whereas the neighboring spheres along the chain can overlap when a is larger than $1.9\text{\AA}$. Non-consecutive spheres interact via a pairwise square-well potential equal




FIG. 4: Dynamical switching between the single helix and the dual helix. (a) A long Monte Carlo trajectory, with standard pivot and crankshaft move sets²⁷, at $T = 0.5\epsilon/k_B$, shows frequent switches between the single helix (lower contact order) and the dual helix (higher contact order). The chain studied here is the same as in Fig. 3. Both the single and the dual helix have similar energies. (b) Similar to panel a but for a much smaller window of time steps and with higher resolution showing a single switch from the single helix to the dual helix. The bottom figures are the snapshots during the switch at times t (in units of 10^6 steps) as indicated.

to $-\epsilon$ within a contact range, R_c . A Frenet frame of reference is assigned to each sphere, i , except to those at the ends of the chain. The tangent vector, $\hat{\mathbf{t}}_i$, in this frame is a unit vector tangential to the circle passing through the centers of beads $i-1$, i and $i+1$. The Frenet normal vector, $\hat{\mathbf{n}}_i$, is a unit vector pointing to the center of this circle. The third vector of the frame is the binormal vector, denoted by $\hat{\mathbf{b}}_i$, and is defined to be the cross product of the tangent and the normal vectors. A side-sphere of radius b is attached to each backbone sphere in direction opposite to the normal with the distance between the centers of the backbone sphere and its side-sphere partner equal to $a+b$. The backbone spheres at the two ends of the chain do not have any side-spheres attached to them. The role of the side-spheres is entirely steric - they are not allowed to overlap with any of the other spheres in the system. The energy of the chain is given

conformation can be written as

$$E = -\varepsilon \sum_{i=1}^{N-2} \sum_{j=i+2}^N \Theta(R_c - |\vec{r}_i - \vec{r}_j|), \quad (1)$$

where $\vec{r}_i$ are center positions of the backbone spheres, N is the number of such spheres, and the step function $\Theta(x)$ is equal to 1 if $x > 0$ and 0 otherwise.

We employ a parallel tempering²⁶ Monte Carlo (MC) scheme for obtaining the ground state as well as other equilibrium characteristics of the system. The simulation entails monitoring 20-30 replicas, each evolving at its own selected temperature, T_i . For each replica, the simulation is carried out with the standard pivot and crankshaft move sets²⁷ and the Metropolis algorithm for move acceptance. In a pivot move, one randomly chooses a sphere i in the chain and rotates the shorter part of the chain (either from 1 through $i-1$ or from $i+1$ through N , where N is the number of spheres in the chain) by a small angle and about a randomly chosen axis that goes through the i -th sphere. In a crankshaft move, two spheres i and j are chosen randomly such that $|i-j| < 6$ and the spheres between i and j are rotated by a small angle and about the axis that goes through i and j . In both move sets, the rotation angle is randomly drawn from a Gaussian distribution of zero mean and a dispersion of 4° . An attempt to exchange replicas i and j is made every 100 MC steps. The exchange is accepted with a probability equal to $p_{ij} = \min\{1, \exp[k_B^{-1}(T_i^{-1} - T_j^{-1})(E_i - E_j)]\}$, where k_B is the Boltzmann constant, and E_i and E_j are the energies of the replicas at the time of the exchange. The weighted multiple histogram technique²⁵ is used to compute the specific heat and the effective free energy.

Appendix: Protein-like folds are adopted by a host of non-biological polymers

Poly(diacetylene)s can form multiple-helical superstructures¹⁵ such as double-helical ribbons.

Poly(ethylene glycol) (PEG) has been shown to assume a helical conformation in isobutyric acid with a trace amount of water without which PEG forms a coil configuration^{16,17}. PEG also forms helices in isopentanoic and n-propanoic acids but not in isobutanol or n-butanol¹⁷. In isobutyric acid, PEG forms a mixture of helices and coils whereas a similar polymer poly(ethylene imine) (PEI) merely forms helices. Phenylacetylene oligomers of specific chain lengths (up to 18 units) and containing a tri(ethylene oxide) side-chain segment at each repeat unit have been found to undergo sharp switching between arrays of random coils and arrays of helical conformations on changing the solvent composition¹⁸. Another interesting example is provided by poly(ethylene oxide) (PEO) dissolved in an electrolyte consisting of a lithium salt¹⁹, LiCF_3SO_3 . In the crystalline phase of the system, the PEO chains are helical and form parallel arrays. In the amorphous phase, the arrays dissolve into separate helices. Finally, there are several examples of synthesis of helical polymers that are discussed by Sanda et al.²⁰ and which include polychloral, polyisocyanates, polyisocyanides, polysilanes, and polycetylenes. Such helical structures can be made to have adjustable geometry and then used to generate nanocavities of tunable sizes²¹. Sanda et al.²⁰ have recognized that the crucial factor for the synthetic polymers to adopt the helical structures is the steric repulsion between the side chains combined with attraction that is usually provided by hydrogen bonding.

Acknowledgments

This work was supported by PRIN 2007, NAFOSTED, the grant N N202 0852 33 from the Ministry of Science and Higher Education in Poland, and the Vietnam Education Foundation (VEF). T.X.H. is a VEF Visiting Scholar.

¹ Katsonis N *et al.* (2006) Reversible conductance switching of single diarylethenes on a gold surface. *Adv. Mater.* 18:13971400.

² Choi B-Y *et al.* (2006) Conformational molecular switch of the azobenzene molecule: A scanning tunnelling microscopy study. *Phys. Rev. Lett.* 96:156106.

³ Liljeroth P, Repp J, Meyer G (2007) Current-induced hydrogen tautomerization and conductance switching of naphthalocyanine molecules. *Science* 317:12031206.

⁴ Dri C, Peters MV, Schwarz J, Hecht S, Grill L (2008) Spatial periodicity in molecular switching, *Nature Nanotechnology* 3:649-653.

⁵ Whitesides GM (2001) The once and future nanomachine. *Scientific American* 285:78-83.

⁶ Sytina OA *et al.* (2008) Conformational changes in an ul-

trafast light-driven enzyme determine catalytic activity. *Nature* 456:1001-1004.

⁷ Schrödinger E (1994) *What is life? The physical aspect of the living cell* (Cambridge University Press, Cambridge).

⁸ Chaikin PM, Lubensky TC (2000) *Principles of condensed matter physics* (Cambridge University Press, Cambridge).

⁹ Rader AJ, Hespenheide BM, Kuhn LA, Thorpe MF (2002) Protein unfolding: Rigidity lost. *Proc. Natl. Acad. Sci. USA* 99:3540.

¹⁰ Szpiro GG (2003) *Kepler's conjecture* (John Wiley & Sons, New York).

¹¹ Zamyatin AA (1972) Protein volume in solution. *Prog. Biophys. Mol. Biol.* 24:107-123.

¹² Banavar JR, Maritan A (2007) Physics of proteins. *Annu. Rev. Biophys. Bio. Struct.* 36:261-280.

- ¹³ Clementi C, Maritan A, Banavar JR (1998) Folding, design, and determination of interaction potentials using off-lattice dynamics of model heteropolymers. *Phys. Rev. Lett.* 81:3287-3290.
- ¹⁴ Magee JE, Vasquez VR, Lue L (2006) Helical structures from an isotropic homopolymer model. *Phys. Rev. Lett.* 96:207802.
- ¹⁵ Weiss J, Jahnke E, Severin N, Rabe JP, Frauenrath H (2008) Consecutive conformational transitions and deaggregations of multiple-helical poly(diacetylene)s. *Nano Lett.* 8:1660-1666.
- ¹⁶ Castellanos P, Norman AI, Greer SC (2006) Conformational change of poly(ethylene glycol) near the critical point of isobutyric acid + water. *J. Phys. Chem. B* 110:22172-22177.
- ¹⁷ Norman AI, Fei Y, Ho DL, Greer SC (2007) Folding and unfolding of polymer helices in solution. *Macromolecules* 40:2559-2567.
- ¹⁸ Wooley KL, Moore JS, Wu C, Yang Y (2000) Novel polymers: molecular to nanoscale order in three dimensions. *Proc. Natl. Acad. Sci. USA* 97:11147-11148.
- ¹⁹ Frech R, Chintapalli S, Bruce PG, Vincent CA (1997) Structure of an amorphous polymer electrolyte, poly(ethylene oxide)₃:LiCF₃SO₃. *Chem. Commun.* 157-158.
- ²⁰ Sanda F, Tabei J, Shiotsuki M, Masuda T (2006) Chiroptical study and conformation analysis of helical polymers surrounded by helical hydrogen-bonding strands. *Science and Technology Advanced Materials* 7:572-577.
- ²¹ Gong B *et al.* (2002) Creating nanocavities of tunable sizes: hollow helices. *Proc. Natl. Acad. Sci. USA* 99:11583-11588.
- ²² Chothia C, Finkelstein AV (1990) The classification and origins of protein folding patterns. *Annu. Rev. Biochem.* 59:1007-1039.
- ²³ Chothia C (1992) One thousand families for the molecular biologist. *Nature* 357:543-544.
- ²⁴ Hoang TX, Trovato A, Seno F, Banavar JR, Maritan A (2004) Geometry and symmetry prescript the free energy landscape of proteins. *Proc. Natl. Acad. Sci. USA* 101:7960-7964.
- ²⁵ Ferrenberg AM, Swendsen RH (1989) Optimized Monte Carlo data analysis. *Phys. Rev. Lett.* 63:1195-1198.
- ²⁶ Marinari E, Parisi G (1992) Simulated tempering: a new Monte Carlo scheme. *Europhys. Lett.* 19:451-458.
- ²⁷ Sokal AD (1996) Monte Carlo methods for self-avoiding walk. *Nucl. Phys. B Proc. Suppl.* 47:172179.

Supporting Information




Figure S1: Correlation functions for a chain of non-overlapping spheres. The tangent-tangent, normal-normal, and the binormal-binormal correlations as a function of sequence separation are shown for a chain of $N = 50$ non-overlapping spheres ($a = 1.9\text{\AA}$) with $R_c = 6.5\text{\AA}$. The chain is studied at $T = 0.6\epsilon/k_B$ which is below the collapse transition. Panel **a** corresponds to the situation without the side-spheres ($b = 0$) and panel **b** with side-spheres ($b = 2.5\text{\AA}$). The tangent-tangent correlation for a sequence separation s is defined as $\langle \hat{\mathbf{t}}_i \cdot \hat{\mathbf{t}}_{i+s} \rangle$ where the average $\langle \cdot \rangle$ is taken over all i running from 2 to $(N - s - 1)$ and over all sample conformations. The normal-normal and binormal-binormal correlations are defined similarly. The correlations are averaged over a run of 10^9 MC steps, in which conformations are considered every 10^5 steps. The conformations displayed at the bottom are typical snapshots obtained in the simulation at two different temperatures as indicated. The two conformations on the left are for $b = 0$ and the remaining two are for $b = 2.5\text{\AA}$. Note that the presence of the side-spheres induce helical conformations and yield qualitatively distinct behaviors of the correlation functions of the tangent, normal, and binormal vectors along the chain.




Figure S2: The crossover between the single helix and the random coil phases in a chain of $N = 50$ overlapping spheres. The parameters are $a = 2.5\text{\AA}$, $b = 2.5\text{\AA}$, and $R_c = 6.5\text{\AA}$. The ground state for this system is a single helix. (a) Dependence of the specific heat on temperature showing a peak at $T_{max} \approx 0.72\epsilon/k_B$. (b) Dependence of the tangent-tangent correlation on the sequence separation for three different temperatures: well-above the transition ($T = 1.6\epsilon/k_B$), just above the transition ($T = 0.8\epsilon/k_B$) and just below the transition ($T = 0.6\epsilon/k_B$). (c) Snapshots of typical conformations at the three temperatures. The specific heat is obtained from fluctuations in the energy using the equation $C = (\langle E^2 \rangle - \langle E \rangle^2)/(k_B T^2)$, where E is the energy and $\langle \cdot \rangle$ indicates an average. Note that the signature of the helical ground state is present in the correlation functions even at a temperature distinctly higher than T_{max} showing that information about the ordered low temperature state is encoded even in the high temperature phase.




Figure S3: Ground state conformations of a chain of $N = 16$ overlapping spheres with side-spheres and an

additional bending energy along the chain. The parameters are $a = 2.5\text{\AA}$, $b = 2.5\text{\AA}$ and $R_c = 6.5\text{\AA}$. The bending energy is given by $\sum_{i=2}^{N-1} e_R(1 - \cos(\theta_i))$, where e_R is a coefficient and θ_i is the angle between two consecutive connectivity vectors $\mathbf{r}_{i-1,i} \equiv \mathbf{r}_i - \mathbf{r}_{i-1}$ and $\mathbf{r}_{i,i+1} \equiv \mathbf{r}_{i+1} - \mathbf{r}_i$ associated with the sphere i . The conformations shown are for $e_R = 2\epsilon$ (a), 4ϵ (b) and 8ϵ (c), respectively. Note that for the largest value of e_R , one obtains a planar hairpin-like conformation.